arXiv:1210.0180v1 [math.OC] 30 Sep 2012

Canonical duality theory and Triality for solving general nonconstrained global optimization problems

D. M. MORALES SILVA[a,1], D. Y. GAO[a,1,*]

[a]*School of Science, Information Technology and Engineering*
University of Ballarat, Victoria 3353

Abstract

General nonconvex optimization problems are studied by using the canonical duality theory. The triality theory is proved for sums of exponentials and quartic polynomials, which solved an open problem left in 2003. This theory can be used to find the global minimum and local extrema. Detailed applications are illustrated by several examples.

Keywords: Canonical duality, triality theory, nonlinear analysis, nonconvex optimization

---

1. Introduction

In this paper, we intend to solve the following nonconvex optimization problem (($\mathcal{P}$) in short):

$$(\mathcal{P}) : \text{ext} \left\{ \Pi(\mathbf{x}) = W(\mathbf{x}) + \frac{1}{2}\mathbf{x}^t \mathbf{A}\mathbf{x} - \mathbf{f}^t\mathbf{x} \mid \mathbf{x} \in \mathbb{R}^n \right\}, \tag{1}$$

where ext$\{*\}$ denotes finding extremum points of a function given in $\{*\}$, $\mathbf{f} \in \mathbb{R}^n$ is a given (input) vector, $\mathbf{A} \in \mathbb{R}^{n\times n}$ is a given symmetric matrix, and $W : \mathbb{R}^n \to \mathbb{R}$ is a combination of fourth order polynomials (double-well functions) and quadratic-exponential functions, namely:

$$W(\mathbf{x}) := \sum_{i\in I_m} \exp\left(\frac{1}{2}\mathbf{x}^t\mathbf{B}_i\mathbf{x} - \alpha_i\right) + \sum_{j\in I_p} \frac{b_j}{2}\left(\frac{1}{2}\mathbf{x}^t\mathbf{C}_j\mathbf{x} - \theta_j\right)^2,$$

---

[*] Corresponding author

Email addresses: `d.moralessilva@ballarat.edu.au` (D. M. MORALES SILVA), `d.gao@ballarat.edu.au` (D. Y. GAO)

[1] FAX: +61 3 5327 9077

where $I_m = \{1, \ldots, m\}, \ I_p = \{1, \ldots, p\}$ are two integer sets with m, p are fixed integers; all the coefficients b_j with $j \in I_p$ are positive constants, and $\alpha_i, \theta_j \in \mathbb{R} \ \ \forall i \in I_m, \ j \in I_p$ are given parameters; the matrices $\{\mathbf{B}_i\}_{i \in I_m}$ and $\{\mathbf{C}_j\}_{j \in I_p}$ are assumed to be symmetric, positive semi-definite such that the cone generated by them contains a positive definite matrix.

The sums of exponential and quartic polynomials $W(\mathbf{x})$ can be used to model a large class of nonlinear phenomena in mathematical physics, engineering mechanics, material sciences, network communication, computational biology, and much more (see [7, 8, 9, 12, 17, 20]). Due to the nonconvexity of the problem, the function Π may possess multiple extremal points. To identify both global and local extremal values is a main task in global optimization.

Canonical duality theory and the associated triality for solving general quartic polynomial optimization problems have been studied recently in [11]. An open problem on the double-min duality statement left by Gao 2003 [5] is solved completely. Based on these results, this paper intends to solve the more challenging problem $(\mathcal{P})$. We will show that by the canonical dual transformation, all critical solutions of $(\mathcal{P})$ can be analytically presented in terms of the canonical dual solutions. The extremality of these solutions can be identified by the triality theory. Several solved examples are listed in the last section.

2. Canonical Dual Problem and Analytical Solutions

Following the standard procedure of the canonical dual transformation (cf. e.g., [5]), first we need to choose a geometric operator $\Lambda = (\Lambda_1(\mathbf{x}), \Lambda_2(\mathbf{x})) : \mathbb{R}^n \to \mathbb{R}^{m+p}$, where

$$\begin{aligned} \Lambda_1(\mathbf{x}) &= \left\{ \frac{1}{2}\mathbf{x}^t \mathbf{B}_i \mathbf{x} - \alpha_i \right\} : \ \mathbb{R}^n \to \mathbb{R}^m, \\ \Lambda_2(\mathbf{x}) &= \left\{ \frac{1}{2}\mathbf{x}^t \mathbf{C}_j \mathbf{x} - \theta_j \right\} : \mathbb{R}^n \to \mathbb{R}^p. \end{aligned}$$

Therefore, the nonconvex function $W(\mathbf{x})$ can be written in the following canonical form

$$W(\mathbf{x}) = V(\Lambda(\mathbf{x})) = V_1(\Lambda_1(\mathbf{x})) + V_2(\Lambda_2(\mathbf{x})) \tag{2}$$

with

$$V_1(\boldsymbol{\epsilon}) = \sum_{i\in I_m} \exp(\epsilon_i), \quad V_2(\boldsymbol{\gamma}) = \sum_{j\in I_p} \frac{1}{2} b_j \gamma_j^2. \tag{3}$$

Clearly, the canonical function $V(\boldsymbol{\varepsilon})$ is convex on

$$\mathcal{V}_a = \{\boldsymbol{\varepsilon} = (\boldsymbol{\epsilon}, \boldsymbol{\gamma}) \in \mathbb{R}^{m+p} |\ \epsilon_i \in [-\alpha_i, +\infty), \ \ \gamma_j \in [-\theta_j, +\infty), \ \ \forall i \in I_m, \ j \in I_p\} \tag{4}$$

such that the canonical dual variable $\boldsymbol{\varsigma} = (\boldsymbol{\tau}, \boldsymbol{\sigma})$ of $\boldsymbol{\varepsilon} = (\boldsymbol{\epsilon}, \boldsymbol{\gamma})$ can be uniquely defined by

$$\boldsymbol{\varsigma} = \nabla V(\boldsymbol{\varepsilon}) \Rightarrow \ \ \boldsymbol{\tau} = \nabla V_1(\boldsymbol{\epsilon}) = \{\exp(\epsilon_i)\}, \ \ \boldsymbol{\sigma} = \nabla V_2(\boldsymbol{\gamma}) = \{b_j \gamma_j\}, \tag{5}$$

and on the canonical dual space

$$\mathcal{V}_a^* = \{\boldsymbol{\varsigma} = (\boldsymbol{\tau}, \boldsymbol{\sigma}) \in \mathbb{R}^{m+p} |\ \tau_i \in [\exp(-\alpha_i), \infty), \ \ \sigma_j \in [-b_j\theta_j, \infty), \ \forall i \in I_m, \ j \in I_p\}, \tag{6}$$

the Legendre conjugate of $V(\boldsymbol{\varepsilon})$ can be defined by

$$V^c(\boldsymbol{\varsigma}) = \text{sta}\{\boldsymbol{\varepsilon}^t \boldsymbol{\varsigma} - V(\boldsymbol{\varepsilon}) |\ \boldsymbol{\varepsilon} \in \mathcal{V}_a\} = V_1^c(\boldsymbol{\tau}) + V_2^c(\boldsymbol{\sigma}) \tag{7}$$

where sta$\{*\}$ denotes finding stationary points of the function given in $\{*\}$ and

$$V_1^c(\boldsymbol{\tau}) = \sum_{i\in I_m} (\tau_i \ln \tau_i - \tau_i), \ \ V_2^c(\boldsymbol{\sigma}) = \sum_{j\in I_p} \frac{1}{2b_j} \sigma_j^2.$$

By using the canonical dual transformation $W(\mathbf{x}) = V(\Lambda(\mathbf{x})) = \Lambda(\mathbf{x})^T \boldsymbol{\varsigma} - V^c(\boldsymbol{\varsigma})$, the Gao-Strang *total complementary function* $\Xi : \mathbb{R}^n \times \mathcal{V}_a^* \to \mathbb{R}$ associated with the problem $(\mathcal{P})$ can be given by

$$\begin{aligned} \Xi(\mathbf{x}, \boldsymbol{\varsigma}) &= \langle \Lambda(\mathbf{x}), \boldsymbol{\varsigma} \rangle - V^c(\boldsymbol{\varsigma}) + \frac{1}{2}\mathbf{x}^t \mathbf{A} \mathbf{x} - \mathbf{f}^t \mathbf{x} \\ &= \frac{1}{2}\mathbf{x}^t \mathbf{G}(\boldsymbol{\varsigma}) \mathbf{x} - \alpha^t \boldsymbol{\tau} - \theta^t \boldsymbol{\sigma} - V_1^c(\boldsymbol{\tau}) - V_2^c(\boldsymbol{\sigma}) - \mathbf{f}^t \mathbf{x}, \end{aligned} \tag{8}$$

where

$$\mathbf{G}(\boldsymbol{\varsigma}) = \mathbf{A} + \sum_{i\in I_m} \tau_i \mathbf{B}_i + \sum_{j\in I_p} \sigma_j \mathbf{C}_j. \tag{9}$$

Via this $\Xi(\mathbf{x}, \boldsymbol{\varsigma})$, the canonical dual function $\Pi^d : \mathcal{V}_a^* \to \mathbb{R}$ can be defined by

$$\Pi^d(\boldsymbol{\varsigma}) := \text{sta}\left\{\Xi(\mathbf{x}, \boldsymbol{\varsigma}) |\ \mathbf{x} \in \mathbb{R}^n\right\} = \left\{\Xi\left(\mathbf{x}(\boldsymbol{\varsigma}), \boldsymbol{\varsigma}\right) : \nabla_{\mathbf{x}}\Xi\left(\mathbf{x}(\boldsymbol{\varsigma}), \boldsymbol{\varsigma}\right) = 0\right\}.$$

Notice that $\nabla_{\mathbf{x}}\Xi(\mathbf{x}, \boldsymbol{\varsigma}) = \mathbf{G}(\boldsymbol{\varsigma})\mathbf{x} - \mathbf{f} = 0$ if and only if

$$\mathbf{G}(\boldsymbol{\varsigma})\mathbf{x} = \mathbf{f}. \tag{10}$$

Let $Col(\mathbf{G}(\boldsymbol{\varsigma}))$ be the space generated by the columns of the matrix $\mathbf{G}(\boldsymbol{\varsigma})$. Then, on the dual feasible space

$$\mathcal{S}_a = \left\{\boldsymbol{\varsigma} \in \mathcal{V}_a^* : \mathbf{f} \in Col(\mathbf{G}(\boldsymbol{\varsigma}))\right\},$$

the primal solution $\mathbf{x} = (\mathbf{G}(\boldsymbol{\varsigma}))^{-1}\mathbf{f}$ is well defined (if $\mathbf{G}(\boldsymbol{\varsigma})$ is singular, $(\mathbf{G}(\boldsymbol{\varsigma}))^{-1}$ denotes its pseudo-inverse, see [1], [15] and references therein) and we have $\Pi^d : \mathcal{S}_a \to \mathbb{R}$

$$\Pi^d(\boldsymbol{\varsigma}) \quad = \quad -\frac{1}{2}\mathbf{f}^t(\mathbf{G}(\boldsymbol{\varsigma}))^{-1}\mathbf{f} - V_1^c(\boldsymbol{\tau}) - V_2^c(\boldsymbol{\sigma}) - \alpha^t\boldsymbol{\tau} - \theta^t\boldsymbol{\sigma}. \tag{11}$$

Therefore, the canonical dual problem is proposed in the following form:

$$(\mathcal{P}^d) : \quad \text{ext}\{\Pi^d(\boldsymbol{\varsigma}) : \boldsymbol{\varsigma} \in \mathcal{S}_a\}. \tag{12}$$

By the canonical duality theory, it is not difficult to show that

$$\Pi(\mathbf{x}) = \text{sta}\{\Xi(\mathbf{x}, \boldsymbol{\varsigma}) : \boldsymbol{\varsigma} \in \mathcal{S}_a\} = \Xi(\mathbf{x}, \boldsymbol{\varsigma}(\mathbf{x})), \tag{13}$$

where $\boldsymbol{\varsigma}(\mathbf{x}) = (\boldsymbol{\tau}(\mathbf{x}), \boldsymbol{\sigma}(\mathbf{x}))$ and

$$(\boldsymbol{\tau}(\mathbf{x}))_i = \exp((\Lambda_1(\mathbf{x}))_i),\ i \in I_m,$$

$$(\boldsymbol{\sigma}(\mathbf{x}))_j = b_j(\Lambda_2(\mathbf{x}))_j,\ j \in I_p.$$

According to the general theory presented in [5], we have the following result.

Theorem 1 (Analytical Solutions). *Suppose that for a given $\mathbf{f} \in \mathbb{R}^n$ the canonical dual space $\mathcal{S}_a$ is not empty. If $\overline{\boldsymbol{\varsigma}} \in \mathcal{S}_a$ is a stationary point of Π^d, then*

$$\overline{\mathbf{x}} = (\mathbf{G}(\overline{\boldsymbol{\varsigma}}))^{-1}\mathbf{f} \tag{14}$$

is a stationary point of Π *and*

$$\Pi(\overline{\mathbf{x}}) = \Pi^d(\overline{\boldsymbol{\varsigma}}). \tag{15}$$

Proof:

Let us calculate $\nabla\Pi^d(\boldsymbol{\varsigma})$ and $\nabla^2\Pi^d(\boldsymbol{\varsigma})$. We know that

$$\nabla\Pi^d(\boldsymbol{\varsigma}) = \begin{bmatrix} \nabla_{\boldsymbol{\tau}}\Pi^d(\boldsymbol{\varsigma}) \\ \nabla_{\boldsymbol{\sigma}}\Pi^d(\boldsymbol{\varsigma}) \end{bmatrix} \in \mathbb{R}^{m+p},$$

then

$$(\nabla_{\boldsymbol{\tau}}\Pi^d(\boldsymbol{\varsigma}))_i = \frac{1}{2}\mathbf{f}^t(\mathbf{G}(\boldsymbol{\varsigma}))^{-1}\mathbf{B}_i(\mathbf{G}(\boldsymbol{\varsigma}))^{-1}\mathbf{f} - \ln\tau_i - \alpha_i,\ i \in I_m; \tag{16}$$

$$(\nabla_{\boldsymbol{\sigma}}\Pi^d(\boldsymbol{\varsigma}))_j = \frac{1}{2}\mathbf{f}^t(\mathbf{G}(\boldsymbol{\varsigma}))^{-1}\mathbf{C}_j(\mathbf{G}(\boldsymbol{\varsigma}))^{-1}\mathbf{f} - \frac{\sigma_j}{b_j} - \theta_j,\ j \in I_p. \tag{17}$$

On the other hand,

$$\nabla^2\Pi^d(\boldsymbol{\varsigma}) = \begin{bmatrix} \nabla^2_{\boldsymbol{\tau\tau}}\Pi^d(\boldsymbol{\varsigma}) & \nabla^2_{\boldsymbol{\tau\sigma}}\Pi^d(\boldsymbol{\varsigma}) \\ \nabla^2_{\boldsymbol{\sigma\tau}}\Pi^d(\boldsymbol{\varsigma}) & \nabla^2_{\boldsymbol{\sigma\sigma}}\Pi^d(\boldsymbol{\varsigma}) \end{bmatrix} \in \mathbb{R}^{m+p} \times \mathbb{R}^{m+p},$$

where $\nabla^2_{\boldsymbol{\tau\sigma}}\Pi^d(\boldsymbol{\varsigma}) := (\nabla_{\boldsymbol{\tau}}(\nabla_{\boldsymbol{\sigma}}\Pi^d(\boldsymbol{\varsigma}))^t)$. Let δ_{ij} be the Kronecker's delta. Then

$$\begin{aligned}
(\nabla^2_{\boldsymbol{\tau\tau}}\Pi^d(\boldsymbol{\varsigma}))_{ij} &= -\mathbf{f}^t(\mathbf{G}(\boldsymbol{\varsigma}))^{-1}\mathbf{B}_i(\mathbf{G}(\boldsymbol{\varsigma}))^{-1}\mathbf{B}_j(\mathbf{G}(\boldsymbol{\varsigma}))^{-1}\mathbf{f} - \frac{\delta_{ij}}{\tau_j}, \\
& \quad i, j \in I_m. \\
(\nabla^2_{\boldsymbol{\tau\sigma}}\Pi^d(\boldsymbol{\varsigma}))_{ij} &= -\mathbf{f}^t(\mathbf{G}(\boldsymbol{\varsigma}))^{-1}\mathbf{B}_i(\mathbf{G}(\boldsymbol{\varsigma}))^{-1}\mathbf{C}_j(\mathbf{G}(\boldsymbol{\varsigma}))^{-1}\mathbf{f} \\
& \quad i \in I_m; j \in I_p. \\
(\nabla^2_{\boldsymbol{\sigma\tau}}\Pi^d(\boldsymbol{\varsigma}))_{ij} &= -\mathbf{f}^t(\mathbf{G}(\boldsymbol{\varsigma}))^{-1}\mathbf{C}_i(\mathbf{G}(\boldsymbol{\varsigma}))^{-1}\mathbf{B}_j(\mathbf{G}(\boldsymbol{\varsigma}))^{-1}\mathbf{f} \\
& \quad i \in I_m; j \in I_p. \\
(\nabla^2_{\boldsymbol{\sigma\sigma}}\Pi^d(\boldsymbol{\varsigma}))_{ij} &= -\mathbf{f}^t(\mathbf{G}(\boldsymbol{\varsigma}))^{-1}\mathbf{C}_i(\mathbf{G}(\boldsymbol{\varsigma}))^{-1}\mathbf{C}_j(\mathbf{G}(\boldsymbol{\varsigma}))^{-1}f - \frac{\delta_{ij}}{b_j} \\
& \quad i, j \in I_p.
\end{aligned}$$

By making $\mathbf{x} = (\mathbf{G}(\boldsymbol{\varsigma}))^{-1}\mathbf{f}$ and $\mathbf{F}(\mathbf{x}) \in \mathbb{R}^{n\times(m+p)}$ be
$\mathbf{F}(\mathbf{x}) = [\mathbf{B}_1\mathbf{x}, \ldots, \mathbf{B}_m\mathbf{x}, \mathbf{C}_1\mathbf{x}, \ldots, \mathbf{C}_p\mathbf{x}]$, we have

$$\nabla^2\Pi^d(\boldsymbol{\varsigma}) = -\mathbf{F}(\mathbf{x})^t(\mathbf{G}(\boldsymbol{\varsigma}))^{-1}\mathbf{F}(\mathbf{x}) - \text{Diag}\ \left(\frac{1}{\tau_1}, \ldots, \frac{1}{\tau_m}, \frac{1}{b_1}, \ldots, \frac{1}{b_p}\right). \tag{18}$$

Let $\mathbf{D} = \text{Diag }(\tau_1, \ldots, \tau_m, b_1, \ldots, b_p)$, then $\nabla^2 \Pi^d(\boldsymbol{\varsigma})$ can be written as

$$\nabla^2 \Pi^d(\boldsymbol{\varsigma}) = -\mathbf{F}(\mathbf{x})^t (\mathbf{G}(\boldsymbol{\varsigma}))^{-1} \mathbf{F}(\mathbf{x}) - \mathbf{D}^{-1}. \tag{19}$$

Calculating $\nabla \Pi(\mathbf{x})$ and $\nabla^2 \Pi(\mathbf{x})$, we have respectively

$$\nabla \Pi(\mathbf{x}) = \sum_{i \in I_m} \exp\left(\frac{1}{2}\mathbf{x}^t \mathbf{B}_i \mathbf{x} - \alpha_i\right) \mathbf{B}_i \mathbf{x} + \sum_{j \in I_p} b_j \left(\frac{1}{2}\mathbf{x}^t \mathbf{C}_j \mathbf{x} - \theta_j\right) \mathbf{C}_j \mathbf{x} + \mathbf{A}\mathbf{x} - \mathbf{f}. \tag{20}$$

$$\begin{aligned} \nabla^2 \Pi(\mathbf{x}) \quad = \quad & \mathbf{A} + \sum_{i \in I_m} \exp\left(\frac{1}{2}\mathbf{x}^t \mathbf{B}_i \mathbf{x} - \alpha_i\right) (\mathbf{B}_i \mathbf{x} (\mathbf{B}_i \mathbf{x})^t + \mathbf{B}_i) \\ & + \sum_{j \in I_p} b_j \left(\mathbf{C}_j \mathbf{x} (\mathbf{C}_j \mathbf{x})^t + \left(\frac{1}{2}\mathbf{x}^t \mathbf{C}_j \mathbf{x} - \theta_j\right) \mathbf{C}_j \right). \end{aligned} \tag{21}$$

Since $\overline{\boldsymbol{\varsigma}} = (\overline{\boldsymbol{\tau}}, \overline{\boldsymbol{\sigma}})$ is a stationary point of Π^d then by Equations (16) and (17) we have that

$$(\Lambda_1(\overline{\mathbf{x}}))_i = \ln \overline{\tau}_i, \; i \in I_m; \tag{22}$$

$$(\Lambda_2(\overline{\mathbf{x}}))_j = \frac{\overline{\sigma}_j}{b_j}, \; j \in I_p. \tag{23}$$

Using Equations (22) and (23) in Equation (20), we obtain

$$\nabla \Pi(\overline{\mathbf{x}}) = \mathbf{G}(\overline{\boldsymbol{\varsigma}})\overline{\mathbf{x}} - \mathbf{f} = \mathbf{G}(\overline{\boldsymbol{\varsigma}})(\mathbf{G}(\overline{\boldsymbol{\varsigma}}))^{-1}\mathbf{f} - \mathbf{f} = 0.$$

Notice that Equations (22) and (23) together with Equations (11) and (13) imply that

$$\Pi(\overline{\mathbf{x}}) = \Xi(\overline{\mathbf{x}}, \overline{\boldsymbol{\varsigma}}) = \Xi((\mathbf{G}(\overline{\boldsymbol{\varsigma}}))^{-1}\mathbf{f}, \overline{\boldsymbol{\varsigma}}) = \Pi^d(\overline{\boldsymbol{\varsigma}}). \tag{24}$$

And this finishes the proof. ■

In the next section, we will show that the extremality of some of these solutions can be identified by a refined triality theory.

3. Triality Theory and Perturbation

Before presenting the refined triality theory, we need the following sets

$$\mathcal{S}_a^+ := \{\boldsymbol{\varsigma} \in \mathcal{S}_a : G(\boldsymbol{\varsigma}) \succeq 0\}, \quad \mathcal{S}_a^- := \{\boldsymbol{\varsigma} \in \mathcal{S}_a : G(\boldsymbol{\varsigma}) \prec 0\}.$$

Lemma 1. *Suppose that* $m+p<n$, $\overline{\boldsymbol{\varsigma}} \in \mathcal{S}_a^-$ *is a stationary point and a local minimizer of* Π^d *and* $\mathbf{x} = (\mathbf{G}(\overline{\boldsymbol{\varsigma}}))^{-1}\mathbf{f}$*. Then, there exists a matrix* $\mathbf{L} \in \mathbb{R}^{n\times(m+p)}$ *with Rank* $(\mathbf{L}) = m+p$ *such that*

$$\mathbf{L}^t \nabla^2 \Pi(\overline{\mathbf{x}}) \mathbf{L} \succeq 0. \tag{25}$$

Proof: Since $\overline{\boldsymbol{\varsigma}} \in \mathcal{S}_a^-$ is a local minimizer of Π^d, we have that $\nabla^2 \Pi^d(\overline{\boldsymbol{\varsigma}}) \succeq 0$. It follows from Equation (19) that

$$-\mathbf{F}(\overline{\mathbf{x}})^t(\mathbf{G}(\overline{\boldsymbol{\varsigma}}))^{-1}\mathbf{F}(\overline{\mathbf{x}}) \succeq \mathbf{D}^{-1} \succ 0.$$

Thus, Rank $(\mathbf{F}(\overline{\mathbf{x}})) = m+p$. Since $\overline{\boldsymbol{\varsigma}} \in \mathcal{S}_a^-$ and $\mathbf{F}(\overline{\mathbf{x}})\mathbf{D}\mathbf{F}(\overline{\mathbf{x}})^t \succeq 0$ there exists a nonsingular matrix $\mathbf{T} \in \mathbb{R}^{n\times n}$ such that

$$\mathbf{T}^t \mathbf{G}(\overline{\boldsymbol{\varsigma}}) \mathbf{T} = \text{Diag }(-\lambda_1, \ldots, -\lambda_n) \tag{26}$$

and

$$\mathbf{T}^t \mathbf{F}(\overline{\mathbf{x}})\mathbf{D}\mathbf{F}(\overline{\mathbf{x}})^t \mathbf{T} = \text{Diag }(a_1, \ldots, a_{m_1+m_2}, 0, \ldots, 0), \tag{27}$$

where $\lambda_i > 0$ for every $i = 1, \ldots, n$ and $a_j > 0$ for every $j = 1, \ldots, m+p$ (see [2], [13] and references therein). According to Lemma 3 in the Appendix, we know that there exists orthogonal matrices $\mathbf{U} \in \mathbb{R}^{n\times n}$ and $\mathbf{E} \in \mathbb{R}^{(m+p)\times(m+p)}$ such that

$$\mathbf{T}^t \mathbf{F}(\overline{\mathbf{x}}) \mathbf{D}^{\frac{1}{2}} = \mathbf{U}\mathbf{R}\mathbf{E}, \tag{28}$$

where $\mathbf{R} \in \mathbb{R}^{n\times(m+p)}$ and

$$\mathbf{R}_{ij} = \begin{cases} \sqrt{a_i}, & i = j \text{ and } i = 1, \ldots, m+p \\ 0, & \text{otherwise.} \end{cases}$$

According to the singular value decomposition theory, we know that $\mathbf{U}$ is the identity matrix. Then

$$\begin{aligned} \nabla^2 \Pi^d(\overline{\boldsymbol{\varsigma}}) &= -\mathbf{F}(\overline{\mathbf{x}})^t(\mathbf{G}(\overline{\boldsymbol{\varsigma}}))^{-1}\mathbf{F}(\overline{\mathbf{x}}) - \mathbf{D}^{-1} \\ &= -(\mathbf{F}(\overline{\mathbf{x}})^t\mathbf{T})[\mathbf{T}^t\mathbf{G}(\overline{\boldsymbol{\varsigma}})\mathbf{T}]^{-1}(\mathbf{T}^t\mathbf{F}(\overline{\mathbf{x}})) - \mathbf{D}^{-1} \\ &= -\mathbf{D}^{-\frac{1}{2}}\mathbf{E}^t\mathbf{R}^t\text{Diag}\left(-\frac{1}{\lambda_1}, \ldots, -\frac{1}{\lambda_n}\right)\mathbf{R}\mathbf{E}\mathbf{D}^{-\frac{1}{2}} - \mathbf{D}^{-1} \succeq 0. \end{aligned}$$

Multiplying by $\mathbf{D}^{\frac{1}{2}}$ from the left and the right

$$\mathbf{D}^{\frac{1}{2}}\nabla^2\Pi^d(\overline{\boldsymbol{\varsigma}})\mathbf{D}^{\frac{1}{2}} = -\mathbf{E}^t\mathbf{R}^t\mathrm{Diag}\,\left(-\frac{1}{\lambda_1},\ldots,-\frac{1}{\lambda_n}\right)\mathbf{R}\mathbf{E} - \mathbf{I}_{(m+p)\times(m+p)} \succeq 0. \tag{29}$$

If we multiply the right side of the last equation by $\mathbf{E}$ from the left and $\mathbf{E}^t$ from the right, we have

$$\begin{aligned} 0 &\preceq -\mathbf{R}^t\mathrm{Diag}\,\left(-\frac{1}{\lambda_1},\ldots,-\frac{1}{\lambda_n}\right)\mathbf{R} - \mathbf{I}_{(m+p)\times(m+p)} \\ &\preceq \mathrm{Diag}\,\left(\frac{a_1}{\lambda_1}-1,\ldots,\frac{a_{m+p}}{\lambda_{m+p}}-1\right), \end{aligned}$$

thus $a_i \geq \lambda_i$, for every $i = 1,\ldots,m+p$. On the other hand

$$\begin{aligned} \mathbf{T}^t\nabla^2\Pi(\overline{\mathbf{x}})\mathbf{T} &= \mathbf{T}^t\mathbf{G}(\overline{\boldsymbol{\varsigma}})\mathbf{T} + \mathbf{T}^t\mathbf{F}(\overline{\mathbf{x}})\mathbf{D}\mathbf{F}(\overline{\mathbf{x}})^t\mathbf{T} \\ &= \mathrm{Diag}\,(-\lambda_1,\ldots,-\lambda_n) + \mathrm{Diag}\,(a_1,\ldots,a_{m+p},0,\ldots,0) \\ &= \mathrm{Diag}\,(a_1-\lambda_1,\ldots,a_{m+p}-\lambda_{m+p},-\lambda_{m+p+1},\ldots,-\lambda_n). \end{aligned}$$

Let $\mathbf{J} \in \mathbb{R}^{n\times n}$ defined by

$$J_{ij} = \begin{cases} 1, & i = j \text{ and } i = 1,\ldots,m+p \\ 0, & \text{otherwise.} \end{cases}$$

Then we have

$$\mathbf{J}^t\mathbf{T}^t\nabla^2\Pi(\overline{\mathbf{x}})\mathbf{T}\mathbf{J} = \mathrm{Diag}\,(a_1-\lambda_1,\ldots,a_{m+p}-\lambda_{m+p}) \succeq 0. \tag{30}$$

Let $\mathbf{L} = \mathbf{T}\mathbf{J}$, clearly Rank $(\mathbf{L}) = m+p$ and $\mathbf{L}^t\nabla^2\Pi(\overline{\mathbf{x}})\mathbf{L} \succeq 0$, this completes the proof. ■

In a similar way, we can prove the following lemma.

Lemma 2. *Suppose that $m+p > n$, $\overline{\boldsymbol{\varsigma}} \in \mathcal{S}_a^-$ is a stationary point Π^d and $\mathbf{x} = (\mathbf{G}(\overline{\boldsymbol{\varsigma}}))^{-1}\mathbf{f}$ is a local minimizer of Π. Then, there exists a matrix $\mathbf{Q} \in \mathbb{R}^{(m+p)\times n}$ with Rank $(\mathbf{Q}) = n$ such that*

$$\mathbf{Q}^t\nabla^2\Pi^d(\overline{\boldsymbol{\varsigma}})\mathbf{Q} \succeq 0. \tag{31}$$

Let the $m+p$ column vectors of $\mathbf{L}$ be respectively as $\mathbf{l}_1, \dots, \mathbf{l}_{m+p}$ and the n column vectors of $\mathbf{Q}$ be respectively as $\mathbf{q}_1, \dots, \mathbf{q}_n$. Clearly, $\mathbf{l}_1, \dots, \mathbf{l}_{m+p}$ are $m+p$ independent vectors and $\mathbf{q}_1, \dots, \mathbf{q}_n$ are n independent vectors. Now the subspaces $\mathcal{X}_b$ and $\mathcal{S}_b$ are defined as follows:

$$\mathcal{X}_b = \left\{ \mathbf{x} \in \mathbb{R}^n : \mathbf{x} = \overline{\mathbf{x}} + \sum_{i=1}^{m+p} \upsilon_i \mathbf{l}_i, \{\upsilon_i\}_{i=1}^{m+p} \subset \mathbb{R} \right\}, \tag{32}$$

$$\mathcal{S}_b = \left\{ \boldsymbol{\varsigma} \in \mathbb{R}^{m+p} : \boldsymbol{\varsigma} = \overline{\boldsymbol{\varsigma}} + \sum_{j=1}^{n} \vartheta_j \mathbf{q}_j, \{\vartheta_j\}_{j=1}^{n} \subset \mathbb{R} \right\}. \tag{33}$$

Now we are ready to present the Refined Triality Theory.

Theorem 2 (Triality Theory). *Let* $\overline{\boldsymbol{\varsigma}}$ *be a stationary point of* Π^d *and* $\overline{\mathbf{x}} = (\mathbf{G}(\overline{\boldsymbol{\varsigma}}))^{-1}\mathbf{f}$. *Assume that* $\det(\nabla^2 \Pi(\overline{\mathbf{x}})) \neq 0$.

(i) If $\overline{\boldsymbol{\varsigma}} \in \mathcal{S}_a^+$, *then* $\overline{\boldsymbol{\varsigma}}$ *is a global maximizer of* Π^d *in* $\mathcal{S}_a^+$ *and* $\overline{\mathbf{x}}$ *is a global minimizer of* Π.

(ii) If $\overline{\boldsymbol{\varsigma}} \in \mathcal{S}_a^-$, *then* $\overline{\boldsymbol{\varsigma}}$ *is a local maximizer of* Π^d *in* $\mathcal{S}_a^-$ *if and only if* $\overline{\mathbf{x}}$ *is a local maximizer of* Π.

(iii) If $\overline{\boldsymbol{\varsigma}} \in \mathcal{S}_a^-$ *and*

  a) if $n = m+p$, *then* $\overline{\boldsymbol{\varsigma}}$ *is a local minimizer of* Π^d *if and only if* $\overline{\mathbf{x}}$ *is a local minimizer of* Π, *i.e., there exists respectively neighborhoods* $\mathcal{X}, \mathcal{S} \subset \mathbb{R}^n$ *of* $\overline{\mathbf{x}}$ *and* $\overline{\boldsymbol{\varsigma}}$ *such that*

$$\Pi(\overline{\mathbf{x}}) = \min_{\mathbf{x} \in \mathcal{X}} \Pi(\mathbf{x}) = \min_{\boldsymbol{\varsigma} \in \mathcal{S}} \Pi^d(\boldsymbol{\varsigma}) = \Pi^d(\overline{\boldsymbol{\varsigma}}); \tag{34}$$

  b) if $m+p < n$ *and* $\overline{\boldsymbol{\varsigma}}$ *is a local minimizer of* Π^d, *then* $\overline{\mathbf{x}}$ *is a saddle point of* Π *and there exists respectively neighborhoods* $\mathcal{X}, \mathcal{S} \subset \mathbb{R}^n$ *of* $\overline{\mathbf{x}}$ *and* $\overline{\boldsymbol{\varsigma}}$, *such that*

$$\Pi(\overline{\mathbf{x}}) = \min_{\mathbf{x} \in \mathcal{X} \cap \mathcal{X}_b} \Pi(\mathbf{x}) = \min_{\boldsymbol{\varsigma} \in \mathcal{S}} \Pi^d(\boldsymbol{\varsigma}) = \Pi^d(\overline{\boldsymbol{\varsigma}}); \tag{35}$$

  c) if $n < m+p$ *and* $\overline{\mathbf{x}}$ *is a local minimizer of* Π, *then* $\overline{\boldsymbol{\varsigma}}$ *is a saddle point of* Π^d *and there exists respectively neighborhoods* $\mathcal{X}, \mathcal{S} \subset \mathbb{R}^n$ *of* $\overline{\mathbf{x}}$ *and* $\overline{\boldsymbol{\varsigma}}$ *such that*

$$\Pi(\overline{\mathbf{x}}) = \min_{\mathbf{x} \in \mathcal{X}} \Pi(\mathbf{x}) = \min_{\boldsymbol{\varsigma} \in \mathcal{S} \cap \mathcal{S}_b} \Pi^d(\boldsymbol{\varsigma}) = \Pi^d(\overline{\boldsymbol{\varsigma}}). \tag{36}$$

Proof:

(i) Since $\overline{\boldsymbol{\varsigma}} \in \mathcal{S}_a^+$, from Equation (19) it is not difficult to show that Π^d is strictly concave in $\mathcal{S}_a^+$ and $\Xi(\cdot, \overline{\boldsymbol{\varsigma}})$ is strictly convex in $\mathbb{R}^n$ and therefore $\overline{\boldsymbol{\varsigma}}$ must be a global maximizer of Π^d in $\mathcal{S}_a^+$ and $\overline{\mathbf{x}}$ is a global minimizer of $\Xi(\cdot, \overline{\boldsymbol{\varsigma}})$. By the definition of Ξ given in Equation (8) and the convexity of V, the Fenchel inequality leads to

$$\Xi(\mathbf{x}, \boldsymbol{\varsigma}) \leq \Pi(\mathbf{x}), \ \forall (\mathbf{x}, \boldsymbol{\varsigma}) \in \mathbb{R}^n \times \mathcal{S}_a.$$

Let us assume now that there exists a vector $\mathbf{x}' \in \mathbb{R}^n$ such that $\Pi(\mathbf{x}') < \Pi(\overline{\mathbf{x}})$, then

$$\Pi(\overline{\mathbf{x}}) > \Pi(\mathbf{x}') \geq \Xi(\mathbf{x}', \overline{\boldsymbol{\varsigma}}) \geq \Xi(\overline{\mathbf{x}}, \overline{\boldsymbol{\varsigma}}) = \Pi(\overline{\mathbf{x}}),$$

where the last equality comes from Equation (24). This contradiction proves that $\overline{\mathbf{x}}$ must be a global minimizer of Π.

(ii) Notice first that using Equations (22) and (23) in Equation (21) we have

$$\nabla^2 \Pi(\overline{\mathbf{x}}) = \mathbf{G}(\overline{\boldsymbol{\varsigma}}) + \mathbf{F}(\overline{\mathbf{x}}) \mathbf{D} \mathbf{F}(\overline{\mathbf{x}})^t, \tag{37}$$

where $F(\mathbf{x})$ and D are defined in Equation (19). If $\overline{\boldsymbol{\varsigma}}$ is a local maximizer of Π^d in $\mathcal{S}_a^-$ we must have that $\nabla^2 \Pi^d(\overline{\boldsymbol{\varsigma}}) \preceq 0$, from Equation (19) this is equivalent to

$$\mathbf{D}^{-1} + \mathbf{F}(\overline{\mathbf{x}})^t (\mathbf{G}(\overline{\boldsymbol{\varsigma}}))^{-1} \mathbf{F}(\overline{\mathbf{x}}) \succeq 0. \tag{38}$$

- If $m + p = n$ and $\mathbf{F}$ is invertible, multiplying Equation (38) by $(\mathbf{F}(\overline{\mathbf{x}})^t)^{-1}$ from the left and $(\mathbf{F}(\overline{\mathbf{x}}))^{-1}$ from the right, we have:

  $$(\mathbf{F}(\overline{\mathbf{x}})^t)^{-1} \mathbf{D}^{-1} (\mathbf{F}(\overline{\mathbf{x}}))^{-1} + (\mathbf{G}(\overline{\boldsymbol{\varsigma}}))^{-1} \succeq 0 \tag{39}$$

  this is equivalent to

  $$(\mathbf{F}(\overline{\mathbf{x}})^t)^{-1} \mathbf{D}^{-1} (\mathbf{F}(\overline{\mathbf{x}}))^{-1} \succeq -(\mathbf{G}(\overline{\boldsymbol{\varsigma}}))^{-1} \succ 0,$$

  which in turn is equivalent to (Lemma 4 in the Appendix)

  $$-\mathbf{G}(\overline{\boldsymbol{\varsigma}}) \succeq \mathbf{F}(\overline{\mathbf{x}}) \mathbf{D} \mathbf{F}(\overline{\mathbf{x}})^t \Longleftrightarrow \nabla^2 \Pi(\overline{\mathbf{x}}) \preceq 0.$$

  By assumption $\det(\nabla^2 \Pi(\overline{\mathbf{x}})) \neq 0$, then $\overline{\mathbf{x}}$ is a local maximum of Π.

- If $m+p \neq n$ or $\mathbf{F}$ is not invertible, then by Lemma 3, there exists orthogonal matrices $\mathbf{E} \in \mathbb{R}^{n \times n}$, $\mathbf{K} \in \mathbb{R}^{(m+p)\times(m+p)}$ and a matrix $\mathbf{R} \in \mathbb{R}^{n \times (m+p)}$ such that

  $$R_{ij} = \begin{cases} s_i, & i = j \text{ and } i = 1, \ldots, r \\ 0, & \text{otherwise} \end{cases}$$

  where $s_i > 0$ for every i, $r = \text{Rank}\ (\mathbf{F}(\overline{\mathbf{x}}))$ and

  $$\mathbf{F}(\overline{\mathbf{x}})\mathbf{D}^{\frac{1}{2}} = \mathbf{ERK}. \tag{40}$$

  Using Equation (40), Equation (38) can be rewritten as:

  $$\mathbf{D}^{-1} + \mathbf{D}^{-\frac{1}{2}}\mathbf{K}^t\mathbf{R}^t\mathbf{E}^t(\mathbf{G}(\overline{\boldsymbol{\varsigma}}))^{-1}\mathbf{ERKD}^{-\frac{1}{2}} \succeq 0$$

  after multiplying this equation by $\mathbf{KD}^{\frac{1}{2}}$ from the left and $\mathbf{D}^{\frac{1}{2}}\mathbf{K}^t$ from the right, we have

  $$\mathbf{I}_{(m+p)\times(m+p)} + \mathbf{R}^t(\mathbf{E}^t\mathbf{G}(\overline{\boldsymbol{\varsigma}})\mathbf{E})^{-1}\mathbf{R} \succeq 0.$$

  This equation is equivalent to

  $$-\mathbf{I}_{(m+p)\times(m+p)} - \mathbf{R}^t(\mathbf{E}^t\mathbf{G}(\overline{\boldsymbol{\varsigma}})\mathbf{E})^{-1}\mathbf{R} \preceq 0.$$

  By Lemma 5 in the Appendix, the last equation is equivalent to

  $$0 \succeq \mathbf{E}^t\mathbf{G}(\overline{\boldsymbol{\varsigma}})\mathbf{E} + \mathbf{RR}^t = \mathbf{E}^t\mathbf{G}(\overline{\boldsymbol{\varsigma}})\mathbf{E} + \mathbf{R}(\mathbf{KD}^{-\frac{1}{2}}\mathbf{DD}^{-\frac{1}{2}}\mathbf{K}^t)\mathbf{R}^t$$

  multiplying by $\mathbf{E}$ from the left and $\mathbf{E}^t$ from the right

  $$0 \succeq \mathbf{G}(\overline{\boldsymbol{\varsigma}}) + (\mathbf{ERKD}^{-\frac{1}{2}})\mathbf{D}(\mathbf{D}^{-\frac{1}{2}}\mathbf{K}^t\mathbf{R}^t\mathbf{E}^t) = \mathbf{G}(\overline{\boldsymbol{\varsigma}}) + \mathbf{F}(\overline{\mathbf{x}})\mathbf{DF}(\overline{\mathbf{x}})^t = \nabla^2\Pi(\overline{\mathbf{x}}).$$

  By the assumption $\det(\nabla^2\Pi(\overline{x})) \neq 0$, $\overline{\mathbf{x}}$ is a local maximum of Π.

Notice that every step of the proof are equivalent, so if $\overline{\mathbf{x}}$ is a local maximum of Π then $\overline{\boldsymbol{\varsigma}}$ must be a local maximum of Π^d.

(iii) Let us consider the three cases:

a) $n = m + p$: if $\overline{\boldsymbol{\varsigma}}$ is a local minimizer of Π^d then

$$\nabla^2 \Pi^d(\overline{\boldsymbol{\varsigma}}) = -\mathbf{F}(\overline{\mathbf{x}})^t (\mathbf{G}(\overline{\boldsymbol{\varsigma}}))^{-1} \mathbf{F}(\overline{\mathbf{x}}) - \mathbf{D}^{-1} \succeq 0$$

$$\Longleftrightarrow -\mathbf{F}(\overline{\mathbf{x}})^t (\mathbf{G}(\overline{\boldsymbol{\varsigma}}))^{-1} \mathbf{F}(\overline{\mathbf{x}}) \succeq \mathbf{D}^{-1}.$$

This implies that Rank $(\mathbf{F}(\overline{\mathbf{x}})) = n$. By multiplying the last inequality by $(\mathbf{F}(\overline{\mathbf{x}})^t)^{-1}$ from the left and by $(\mathbf{F}(\overline{\mathbf{x}}))^{-1}$ from the right, we have

$$-(\mathbf{G}(\overline{\boldsymbol{\varsigma}}))^{-1} \succeq (\mathbf{F}(\overline{\mathbf{x}})^t)^{-1} \mathbf{D}^{-1} (\mathbf{F}(\overline{\mathbf{x}}))^{-1}.$$

By Lemma 4 this is equivalent to

$$-\mathbf{G}(\overline{\boldsymbol{\varsigma}}) \preceq \mathbf{F}(\overline{\mathbf{x}}) \mathbf{D} \mathbf{F}(\overline{\mathbf{x}})^t \Longleftrightarrow \nabla^2 \Pi(\overline{\mathbf{x}}) \succeq 0.$$

And since $\det(\nabla^2 \Pi(\overline{x})) \neq 0$, $\overline{\mathbf{x}}$ is a local minimizer of Π. In a similar way we can prove the converse.

b) From Equation (19) we know that

$$-\mathbf{F}(\overline{\mathbf{x}})^t (\mathbf{G}(\overline{\boldsymbol{\varsigma}}))^{-1} \mathbf{F}(\overline{\mathbf{x}}) \succeq \mathbf{D}^{-1},$$

then $-\mathbf{F}(\overline{\mathbf{x}})^t (\mathbf{G}(\overline{\boldsymbol{\varsigma}}))^{-1} \mathbf{F}(\overline{\mathbf{x}})$ is a nonsingular matrix and Rank $(\mathbf{F}(\overline{\mathbf{x}})) = m + p < n$. We claim now that $\overline{\mathbf{x}}$ is not a local minimizer of Π. This is because that if $\overline{\mathbf{x}}$ is also a local minimizer, we would have

$$\nabla^2 \Pi(\overline{\mathbf{x}}) = \mathbf{G}(\overline{\boldsymbol{\varsigma}}) + \mathbf{F}(\overline{\mathbf{x}}) \mathbf{D} \mathbf{F}(\overline{\mathbf{x}})^t \succeq 0,$$

thus

$$\mathbf{F}(\overline{\mathbf{x}}) \mathbf{D} \mathbf{F}(\overline{\mathbf{x}})^t \succeq -\mathbf{G}(\overline{\boldsymbol{\varsigma}}).$$

This implies that

$$n = \text{Rank } (-\mathbf{G}(\overline{\boldsymbol{\varsigma}})) = \text{Rank } (\mathbf{F}(\overline{\mathbf{x}}) \mathbf{D} \mathbf{F}(\overline{x})^t) = m + p,$$

which is a contradiction. Therefore, $\overline{\mathbf{x}}$ is a saddle point of Π.

To prove Equation (35), we let L be as in Lemma 1 and $\{\mathbf{l}_i\}_{i=1}^{m+p}$ be the column vectors of L. Define

$$\varphi(t_1, \ldots, t_{m+p}) := \Pi(\overline{\mathbf{x}} + t_1 \mathbf{l}_1 + \ldots + t_{m+p} \mathbf{l}_{m+p}).$$

We need to show that $(0,\dots,0)\in\mathbb{R}^{m+p}$ is a local minimizer of the function φ. Notice that

$$\nabla\varphi(0,\dots,0)=\mathbf{L}^t\nabla\Pi(\overline{\mathbf{x}})=0$$

and

$$\nabla^2\varphi(0,\dots,0)=\mathbf{L}^t\nabla^2\Pi(\overline{\mathbf{x}})\mathbf{L}\succeq 0,$$

which is a consequence of Lemma 1. Furthermore, from Equation (30) we have that

$$\nabla^2\varphi(0,\dots,0)=\text{Diag }(a_1-\lambda_1,\dots,a_{m+p}-\lambda_{m+p}),$$

and since $\det(\nabla^2\Pi(\overline{\mathbf{x}}))\neq 0$ it can be proven that $a_i>\lambda_i$ for every i. The proof is complete.

c) The proof is similar to item b). ■

Remark: The triality theory states precisely that if $\boldsymbol{\varsigma}$ is a global maximizer of Π^d on a certain set, then $\mathbf{x}$ is a global minimizer for Π. This is known from the general result by Gao and Strang in [10]. If $\boldsymbol{\varsigma}$ is a local maximizer for Π^d then $\mathbf{x}$ is also a local maximizer for Π. This is the so-called double-max duality statement. If $\boldsymbol{\varsigma}$ is a local minimizer for Π^d, then $\mathbf{x}$ is also a local minimizer for Π in certain directions. This is so-called double-min duality in the standard triality form proposed in [4]. The triality theory was first discovered in nonconvex mechanics [3]. It was realized in 2003 that the double-min duality holds under certain additional condition, which was left as an open problem (see [5, 6]). Recently, this open problem is solved for quartic polynomial optimization problem [11]. This result is now generalized to the general nonconvex problem $(\mathcal{P})$. Part (iii) of Theorem 2 shows that if $m+p=n$, then $\boldsymbol{\varsigma}$ is a local minimizer if and only if $\mathbf{x}$ is also a local minimizer. In other cases either $\mathbf{x}$ is a saddle point of Π or $\boldsymbol{\varsigma}$ is a saddle point of Π^d.

4. Numerical Examples

In the following examples, $m=p=1$ and $b_1=1$. The graphs provided and the numerical results were obtained using Maxima [14].



(a) Contour Levels of Π.

(b) Graph of Π.

Figure 1: Π function of Example 1

4.1. Example 1: Unique solution

First, we consider the case that the primal function has a unique critical point. We let $\alpha_1 = \theta_1 = 1$ and

$$\mathbf{A} = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}, \mathbf{B}_1 = \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix}, \mathbf{C}_1 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \mathbf{f} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}.$$

Clearly, the function $\Pi : \mathbb{R}^2 \to \mathbb{R}$ is given by

$$\Pi(x, y) = \exp\left(\frac{1}{2}(x^2 + 2y^2) - 1\right) + \frac{1}{2}\left(\frac{1}{2}(x^2 + y^2) - 1\right)^2 + \frac{1}{2}(x^2 - y^2) - x - y,$$

and the dual function has the form of

$$\Pi^d(\tau, \sigma) = -\frac{1}{2}\left(\frac{1}{1 + \tau + \sigma} + \frac{1}{2\tau + \sigma - 1}\right) - \tau \cdot \ln(\tau) - \frac{1}{2}\sigma^2 - \sigma.$$

It can be shown that Π^d has only one critical point in $\mathcal{S}_a^+$ and it is given (approximately) by

$$\overline{\boldsymbol{\varsigma}} = (1.171057661103504, -0.34599084656216).$$

By the triality theory, the vector

$$\overline{\mathbf{x}} = \mathbf{G}(\overline{\boldsymbol{\varsigma}})^{-1}\mathbf{f} = (0.54792514555217, 1.003890602479819)$$

is a global minimizer of the primal problem.



(a) Contour Levels of Π^d.

(b) Graph of Π^d.

Figure 2: Π^d function of Example 1

4.2. Example 2: Two solutions

We now consider the case that the primal function has two critical points. Let $\alpha_1 = 1,\ \ \theta_1 = 50$, and

$$\mathbf{A} = \begin{bmatrix} 1 & 0 \\ 0 & -16 \end{bmatrix}, \mathbf{B}_1 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \mathbf{C}_1 = \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix}, \mathbf{f} = \begin{bmatrix} -25 \\ 9 \end{bmatrix}.$$

The primal function $\Pi : \mathbb{R}^2 \to \mathbb{R}$ is then given by

$$\Pi(x, y) = \exp\left(\frac{1}{2}(x^2 + y^2) - 1\right) + \frac{1}{2}\left(\frac{1}{2}(x^2 + 2y^2) - 50\right)^2 + \frac{1}{2}(x^2 - 16y^2) + 25x - 9y$$

and its canonical dual is

$$\Pi^d(\tau, \sigma) = -\frac{1}{2}\left(\frac{81}{-16 + \tau + 2\sigma} + \frac{625}{1 + \tau + \sigma}\right) - \tau \cdot \ln(\tau) - \frac{1}{2}\sigma^2 - 50\sigma,$$

which has two critical points:

$$\overline{\varsigma_1} = (96.61711963278241, -38.94928057661689) \in \mathcal{S}_a^+,$$

$$\overline{\varsigma_2} = (0.42157060067968, -49.86072154366873) \in \mathcal{S}_a^-.$$

Therefore, by the triality theory, the associated vector

$$\overline{\mathbf{x}_1} = \mathbf{G}(\overline{\varsigma_1})^{-1}\mathbf{f} = (-0.42612784793499, 3.310578038951848)$$

is a global minimizer of $\Pi(\mathbf{x})$ and

$$\overline{\mathbf{x}_2} = (0.51611144112381, -0.078057328303129)$$

is a local maximizer (see Figure 3) since $\overline{\varsigma_2}$ is a local maximum of Π^d in $\mathcal{S}_a^-$ (see Figure 5).



(a) Contour Levels of Π. (b) Graph of Π.

Figure 3: Primal function Π in Example 2



(a) Contour Levels of Π^d in $\mathcal{S}_a^+$. (b) Graph of Π^d in $\mathcal{S}_a^+$.

Figure 4: Π^d function in $\mathcal{S}_a^+$ of Example 2



(a) Contour Levels of Π^d in $\mathcal{S}_a^-$. (b) Graph of Π^d in $\mathcal{S}_a^-$.

Figure 5: Π^d function in $\mathcal{S}_a^-$ of Example 2



(a) Contour Levels of Π. (b) Graph of Π.

Figure 6: Π function of Example 3

4.3. Example 3: Multi-solutions

In order to illustrate the triality theory, we let $\alpha_1 = \theta_1 = 2$, and

$$\mathbf{A} = \begin{bmatrix} -16 & 0 \\ 0 & -4 \end{bmatrix}, \mathbf{B}_1 = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \mathbf{C}_1 = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}, \mathbf{f} = \begin{bmatrix} 2 \\ 2 \end{bmatrix}.$$

Accordingly, we have

$$\Pi(x,y) = \exp\left(\frac{1}{2}x^2 - 2\right) + \frac{1}{2}\left(\frac{1}{2}y^2 - 2\right)^2 + \frac{1}{2}(-16x^2 - 4y^2) - 2x - 2y,$$

$$\Pi^d(\tau,\sigma) = -\frac{1}{2}\left(\frac{4}{\sigma - 4} + \frac{4}{\tau - 16}\right) - \tau \cdot \ln(\tau) - \tau - \frac{1}{2}\sigma^2 - 2\sigma.$$

In this case, Π^d has total six critical points but only one

$$\overline{\varsigma_1} = (16.64468576727409, 4.552474610531074) \in \mathcal{S}_a^+,$$

(see Figure 7) and two

$$\overline{\varsigma_2} = (0.13641513779858, -1.943380912562619) \in \mathcal{S}_a^-,$$

$$\overline{\varsigma_3} = (15.34981976568548, 3.390906302031545) \in \mathcal{S}_a^-.$$

From Figures 8 we can see that $\overline{\varsigma_2}$ is a local maximizer and $\overline{\varsigma_3}$ is a local minimizer of Π^d. Therefore, by the triality theory, we know that

$$\overline{\mathbf{x}_1} = \mathbf{G}(\overline{\varsigma_1})^{-1}\mathbf{f} = (3.102286573591542, 3.620075858467906)$$

is a global minimizer;

$$\overline{\mathbf{x}_2} = (-0.12607490787063, -0.33650880356205)$$



(a) Contour Levels of Π^d in $\mathcal{S}_a^+$. (b) Graph of Π^d in $\mathcal{S}_a^+$.

Figure 7: Π^d function in $\mathcal{S}_a^+$ of Example 3



(a) Contour Levels of Π^d in $\mathcal{S}_a^-$. (b) Graph of Π^d in $\mathcal{S}_a^-$.

Figure 8: Π^d function in $\mathcal{S}_a^-$ of Example 3

is a local maximizer and

$$\overline{\mathbf{x}_3} = (-3.076070133243102, -3.283567054905852)$$

is a local minimizer of $\Pi(\mathbf{x})$ (see Figure 6).

4.4. Example 4: Non-unique global minima

In the case that $\mathcal{S}_a^+$ is empty, the primal problem could have more than one global minima. To see this, we let $\mathbf{f} \equiv 0,\ \alpha_1 = \theta_1 = 2$, and

$$\mathbf{A} \equiv 0, \mathbf{B}_1 = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \mathbf{C}_1 = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}.$$

In this case, the primal function

$$\Pi(x, y) = \exp\left(\frac{1}{2}x^2 - 2\right) + \frac{1}{2}\left(\frac{1}{2}y^2 - 2\right)^2$$

has 2 global minimums at $(0, -2)$, $(0, 2)$ and a local maximum at $(0, 0)$. While the dual function

$$\Pi^d(\tau, \sigma) = -\tau \ln \tau - \tau - \frac{1}{2}\sigma^2 - 2\sigma$$

does not have a stationary point in $\mathcal{S}_a^+$. Indeed, its only critical point is in the boundary of its domain: $\overline{\boldsymbol{\varsigma}} = (\exp(-2), -2)$. By defining $\overline{\mathbf{x}} = \mathbf{G}(\overline{\boldsymbol{\varsigma}})^{-1}\mathbf{f}$, we have that $\overline{\mathbf{x}} = (0, 0)$.

In order to find a global minimum of Π, we need to introduce the following perturbations:

$$\mathbf{A}_n = \begin{bmatrix} -\frac{16}{n} & 0 \\ 0 & -\frac{4}{n} \end{bmatrix} \text{ and } \mathbf{f_n} = \begin{bmatrix} \frac{2}{n} \\ \frac{2}{n} \end{bmatrix}, \text{ for every } n \in I\!N.$$

Then, the associated primal and dual functions are

$$\Pi_n(x,y) = \exp\left(\frac{1}{2}x^2 - 2\right) + \frac{1}{2}\left(\frac{1}{2}y^2 - 2\right)^2 + \frac{1}{2}\left(-\frac{16}{n}x^2 - \frac{4}{n}y^2\right) - \frac{2}{n}x - \frac{2}{n}y,$$

$$\Pi_n^d(\tau,\sigma) = -\frac{1}{2}\left(\frac{4}{n^2\left(\tau - \frac{16}{n}\right)} + \frac{4}{n^2\left(\sigma - \frac{4}{n}\right)}\right) - \tau\ln\tau + \tau - \frac{1}{2}\sigma^2 - 2\tau - 2\sigma.$$

Notice that if $n = 1$ we are in the case presented in Example 3. Let us show that for sufficiently large values of n we can find a stationary point for Π_n^d in $\mathcal{S}_a^+$, namely $\overline{\varsigma_n}$. Furthermore, by defining $\overline{\mathbf{x}}_n = \mathbf{G}(\overline{\boldsymbol{\varsigma}}_n)^{-1}\mathbf{f_n}$ we will have a convergent sequence.

Let us calculate the gradient of Π_n^d:

$$\nabla\Pi_n^d(\tau,\sigma) = \begin{bmatrix} -2 - \ln\tau + \frac{2}{(n\tau-16)^2} \\ -\sigma - 2 + \frac{2}{(n\sigma-4)^2} \end{bmatrix}.$$

Let $h(\tau) = -2 - \ln\tau + \frac{2}{(n\tau-16)^2}$ and $g(\sigma) = -\sigma - 2 + \frac{2}{(n\sigma-4)^2}$. It is not difficult to show that there exists a sufficiently large $N \in I\!N$, such that if $n > N$, the following are true:

a) $n \cdot \exp\left(-2 + \dfrac{1}{n}\right) - 16$ and $n \cdot \exp(-2) - 16$ are positive numbers.

b) $$h\left(\exp\left(-2 + \frac{1}{n}\right)\right) = \frac{2}{(n \cdot \exp(-2 + \frac{1}{n}) - 16)^2} - \frac{1}{n}$$
$$< 0 < h(\exp(-2)) = \frac{2}{(n \cdot \exp(-2) - 16)^2}.$$

c) $g\left(\dfrac{5.1}{n}\right) \approx -\dfrac{5.1}{n} - 0.34710743801 < 0 < g\left(\dfrac{4.9}{n}\right) \approx 0.46913580247 - \dfrac{4.9}{n}.$

Based on these results, we know that for every $n > N$, $\nabla\Pi_n^d$ has a stationary point $\overline{\boldsymbol{\varsigma}}_n = (\overline{\tau}_n, \overline{\sigma}_n) \in [\exp(-2), \exp(-2 + \frac{1}{n})] \times \left[\frac{4.9}{n}, \frac{5.1}{n}\right]$. Moreover, by the fact that $g(\overline{\sigma}_n) = 0$, it is easy to obtain $\lim_{n\to+\infty} n \cdot \overline{\sigma}_n = 5$.

Notice also that

$$\mathbf{G}(\overline{\boldsymbol{\varsigma}}_n) = \begin{bmatrix} \overline{\tau}_n - \dfrac{16}{n} & 0 \\ 0 & \overline{\sigma}_n - \frac{4}{n} \end{bmatrix}$$

is positive definite. Therefore, the perturbed solution can be obtained as

$$\overline{\mathbf{x}}_n = \mathbf{G}(\overline{\boldsymbol{\varsigma}}_n)^{-1}\mathbf{f_n} = \begin{bmatrix} \frac{2}{n\cdot\overline{\tau}_n - 16} \\ \frac{2}{n\cdot\overline{\sigma}_n - 4} \end{bmatrix}.$$

Since $\overline{\tau}_n \in [\exp(-2), \exp(-2 + \frac{1}{n})]$ then $\lim_{n\to+\infty} \tau_n = \exp(-2)$. By the fact that $\lim_{n\to+\infty} n \cdot \overline{\sigma}_n = 5$, we have

$$\lim_{n\to+\infty} \overline{\mathbf{x}}_n = \begin{bmatrix} 0 \\ 2 \end{bmatrix},$$

which is a solution of Π.

Canonical perturbation method was originally introduce in [16] for solving nonconvex polynomial minimization problems. This method has been used successfully in integer programming and network communication (see [9, 19]).

5. Appendix: Some Lemmas in Matrix Analysis

The following results are needed in the proofs of Section 2.

Lemma 3. *(Singular value decomposition [13]) For any given matrix* $\mathbf{M} \subset \mathbb{R}^{m\times n}$ *with Rank* $(\mathbf{M}) = r$, *there exists* $\mathbf{U} \subset \mathbb{R}^{m\times m}$, $\mathbf{R} \subset \mathbb{R}^{m\times n}$ *and* $\mathbf{E} \subset \mathbb{R}^{n\times n}$ *such that*

$$\mathbf{M} = \mathbf{URE};$$

where $\mathbf{U}$ *and* $\mathbf{E}$ *are orthogonal matrices, and*

$$R_{ij} = \begin{cases} s_i, & i = j,\ i = 1, \ldots, r \\ 0, & i \neq j, \end{cases}$$

where $s_i > 0$ *for every* $i = 1, \ldots, r$.

Lemma 4. *[13] If* $\mathbf{G}$ *and* $\mathbf{U}$ *are positive definite matrices in* $\mathbb{R}^{n\times n}$*, then* $\mathbf{G} \succeq \mathbf{U}$ *if and only if* $\mathbf{U}^{-1} \succeq \mathbf{G}^{-1}$.

Lemma 5. *[11] Suppose* $\mathbf{P}$*,* $\mathbf{U}$ *and* $\mathbf{D}$ *are three matrices in* $\mathbb{R}^{n\times n}$ *such that*

$$\mathbf{D} = \begin{bmatrix} \mathbf{D}_{11} & 0_{m\times(n-m)} \\ 0_{(n-m)\times n} & 0_{(n-m)\times(n-m)} \end{bmatrix},$$

where $\mathbf{D}_{11} \in \mathbb{R}^{m\times m}$ *is nonsingular and*

$$\mathbf{P} = \begin{bmatrix} \mathbf{P}_{11} & \mathbf{P}_{12} \\ \mathbf{P}_{21} & \mathbf{P}_{22} \end{bmatrix} \prec 0, \ \mathbf{U} = \begin{bmatrix} \mathbf{U}_{11} & 0_{m\times(n-m)} \\ 0_{(n-m)\times m} & \mathbf{U}_{22} \end{bmatrix} \succ 0,$$

$\mathbf{P}_{ij}$ *and* $\mathbf{U}_{ii}$ *are appropriate dimensional matrices for* $i, j = 1, 2$*. Then,*

$$\mathbf{P} + \mathbf{D}\mathbf{U}\mathbf{D}^t \preceq 0 \Longleftrightarrow -\mathbf{D}^t\mathbf{P}^{-1}\mathbf{D} - \mathbf{U}^{-1} \preceq 0. \tag{41}$$

Acknowledgements This research is supported by US Air Force Office of Scientific Research under the grant AFOSR FA9550-10-1-0487.

References

[1] Desoer, C. A.; Whalen, B. H. *A Note on Pseudoinverses.* Journal of the Society for Industrial and Applied Mathematics, Vol. 11, No 2, pp. 442-447 (1963).

[2] Feng, J. M.; Lin, G. X.; Sheu, R. L.; Xia, Y. *Duality and solutions for quadratic programming over single non-homogeneous quadratic constraint.* Journal of Global Optimization, Published online: 17 Nov 2010.

[3] Gao, D.Y.: *Dual extremum principles in finite deformation theory with applications to post-buckling analysis of extended nonlinear beam theory.* Applied Mechanics Reviews, 50 (11), S64-S71 (1997).

[4] Gao, D. Y. *Duality Principles in nonconvex systems. Theory Methods and Applications.* Kluwer Academic Publishers, Dordrecht/Boston, 2000.

[5] Gao, D.Y. *Perfect duality theory and complete solutions to a class of global optimization problems. Optim.* 52(4–5), pp. 467-493 (2003).

[6] Gao, D.Y.: Nonconvex semi-linear problems and canonical dual solutions. In: Gao, D.Y., Ogden, R.W. (eds.) *Advances in Mechanics and Mathematics,* vol. II, pp. 261312. Kluwer Academic Publishers, Dordrecht (2003)

[7] Gao, D. Y.; Ogden R. W. *Multiple solutions to non-convex variational problems with implications for phase transitions and numerical computation.* Q. J. Mech. Appl. Math, Vol. 61. No. 4 (2008), 497-522.

[8] Gao, D. Y.; Ruan N. *Solutions and optimality criteria for nonconvex quadratic-exponential minimization problem,.* Math. Meth. Operations Research, 67 (3), pp. 479-491 (2008).

[9] Gao, D.Y., Ruan, N, and Pardalos, P.M. Canonical dual solutions to sum of fourth-order polynomials minimization problems with applications to sensor network localization, in *Sensors: Theory, Algorithms and Applications*, P.M. Pardalos, Y.Y. Ye, V. Boginski, and C. Commander (eds). Springer(2010).

[10] Gao, D.Y. and Strang, G. *Geometric nonlinearity: Potential energy, complementary energy, and the gap function.* Quart. Appl. Math. 47 (3), pp. 487-504 (1989).

[11] Gao, D. Y. and Wu, C. *On the triality theory for a quartic polynomial optimization problem,* J. Industrial and Management Optimization, 8(1):229-242, (2012).

[12] Gao, D.Y. and Yu, H.F.: Multi-scale modelling and canonical dual finite element method in phase transitions of solids. *Int. J. Solids and Structures*, 45:3660–3673 (2008).

[13] Horn, R. A.; Johnson, C. R. *Matrix Analysis.* Cambridge University Press (1985).

[14] Maxima.sourceforge.net. *Maxima, a Computer Algebra System.* Version 5.22.1 (2010). http://maxima.sourceforge.net/

[15] Peters, G.; Wilkinson J. H. *The least squares problem and pseudo-inverses.* The Computer Journal, Vol. 13, No 3, pp. 309-316 (1970).

[16] Ruan, N., Gao, D.Y., and Jiao, Y.: Canonical dual least square method for solving general nonlinear systems of quadratic equations, *Computational Optimization and Applications*, Vol 47, 335-347 (2010).

[17] Santos, H.A.F.A. and Gao D.Y.: Canonical dual finite element method for solving post-buckling problems of a large deformation elastic beam. *Int. J. Nonlinear Mechanics,* 2011, doi:10.1016/j.ijnonlinmec.2011.05.012

[18] Sewell, M.J. *Maximum and minimum principles.* Cambridge University Press, Cambridge, New York, Port Chester, Melbourne Sydney (1987).

[19] Wang, Z.B., Fang, S.C., Gao, D.Y., and Xing, W.X.: Canonical dual approach to solving the maximum cut problem, to appear in *J. Glob. Optim.*

[20] Zhang J., Gao, D.Y. and Yearwood, J.: A novel canonical dual computational approach for prion AGAAAAGA amyloid fibril molecular modeling. *Journal of Theoretical Biology*, 284, 149-157 (2011). doi:10.1016/j.jtbi.2011.06.024

[21] Zia, R. K. P.; Redish, E. F.; McKay S. R. *Making Sense of the Legendre Transform* American Journal of Physics, Vol. 77, Issue 7, pp. 614-622 (2009).


-34
-36
-38
-40
-42
-44
-46
-48
-50
-52
-54
-56
-58
5
4
3
2
1
0
-1
2
4
6
8
10
12
14
16
18
20


-34
-36
-38
-40
-42
-44
-46
-48
-50
-52
-54
-56
-58
-30
-35
-40
-45
-50
-55
-60
2
4
6
8
10
12
14
16
18
20
-1
0
1
2
3
4
5